

## 兔抗 NDUFB11 多克隆抗体

中文名称: 兔抗 NDUFB11 多克隆抗体

英文名称: Anti-NDUFB11 rabbit polyclonal antibody

别 名: ESSS; Np15; P17.3; NP17.3; CI-ESSS; LSDMCA3

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗原: NDUFB11

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

Complex 1 (also known as NADH dehydrogenase) of the electron transport chain (respiratory chain) is an enzymat ic complex that catalyzes the transfer of electrons from NADH to ubiquinone. Free energy from the reaction is c onserved in the transfer of protons into the intermembra ne space to create an electrochemical proton gradient, a driving force for ATP synthesis. Complex 1 is a complicat ed, multi-protein, L-shaped complex composed of at leas t 45 different subunits and located in the mitochondrial i nner membrane. NDUFB11 (NADH dehydrogenase (ubiquinone) 1 beta subcomplex subunit 11), also known as ES



	SS, Np15, Np17.3 (neuronal protein 17.3) or p17.3, is a h ydrophobic transmembrane protein belonging to the Co mplex I NDUFB11 subunit family. Ubiquitously expressed, NDUFB11 localizes to the inner membrane of the mitoch ondrion and functions as an accessory subunit of Compl ex I. The cAMP-dependent phosphorylation of NDUFB11 is important for the regulation of Complex I activity.
Applications:	ELISA, WB, IHC
Name of antibody:	NDUFB11
Immunogen:	Full length fusion protein
Full name:	NADH:ubiquinone oxidoreductase subunit B11
Synonyms:	ESSS; Np15; P17.3; NP17.3; CI-ESSS; LSDMCA3
SwissProt:	Q9NX14
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human lung cancer and human prostate cancer
IHC Recommend dilution:	50-300
WB Predicted band size:	17 kDa
WB Positive control:	Mouse brain tissue , Mouse liver tissue , 231 cell , K562 c ell , A431 cell lysates
WB Recommended dilution:	1000-5000









