

## H1-6 抗原(重组蛋白)

- 中文名称: H1-6 抗原 (重组蛋白)
- 英文名称: H1-6 Antigen (Recombinant Protein)
- 别 名: H1.6 linker histone, cluster member; H1t; H1.6; H1FT; HIST1H1T; dJ221C16.2
- 相关类别: 抗原
- 储 存: 冷冻 (-20℃)

## 概述

Fusion protein corresponding to a region derived from 1-207 amino acids of human H1-6

## 技术规格

Full name:	H1.6 linker histone, cluster member	
Synonyms:	H1t; H1.6; H1FT; HIST1H1T; dJ221C16.2	
Swissprot:	P22492	
Gene Accession:	BC130350	
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE	
Expression system:	Escherichia coli	
Tags:	His tag C-Terminus, GST tag N-Terminus	
Background:	Histones are basic nuclear proteins responsible for nucleosome struc ture of the chromosomal fiber in eukaryotes. Two molecules of eac h of the four core histones (H2A, H2B, H3, and H4) form an octam er, around which approximately 146 bp of DNA is wrapped in repea ting units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless an d encodes a replication-dependent histone that is a member of the histone H1 family. Transcripts from this gene lack polyA tails but in	



stead	contain a palindromic termination element. This gene is for	und
in the	e large histone gene cluster on chromosome 6. [provided by	y R
efSeq	J, Aug 2015]	