

## 兔抗 TXNDC12 多克隆抗体

中文名称：兔抗 TXNDC12 多克隆抗体

英文名称： Anti-TXNDC12 rabbit polyclonal antibody

别名： thioredoxin domain containing 12; AG1; AGR1; ERP16; ERP18; ERP19; TLP19; hAG-1; PDIA16; hTLP19

相关类别： 一抗

储存： 冷冻 (-20℃)

宿主： Rabbit

抗原： TXNDC12

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

**Background:**

This gene encodes a member of the thioredoxin superfamily. Members of this family are characterized by a conserved active motif called the thioredoxin fold that catalyzes disulfide bond formation and isomerization. This protein localizes to the endoplasmic reticulum and has a single atypical active motif. The encoded protein is mainly involved in catalyzing native disulfide bond formation and displays activity similar to protein-disulfide isomerases. This protein may play a role in defense against endoplasmic reticulum stress. Alternate splicing results in

	n both coding and non-coding variants.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	TXNDC12
<b>Immunogen:</b>	Fusion protein of human TXNDC12
<b>Full name:</b>	thioredoxin domain containing 12
<b>Synonyms:</b>	AG1; AGR1; ERP16; ERP18; ERP19; TLP19; hAG-1; PDIA16 ; hTLP19
<b>SwissProt:</b>	O95881
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human colorectal cancer and Human liver cancer
<b>IHC Recommend dilution:</b>	50-200



