

兔抗 RELA (Phospho-Thr505)多克隆抗体

中文名称: 兔抗 RELA (Phospho-Thr505)多克隆抗体

英文名称: Anti-RELA (Phospho-Thr505) rabbit polyclonal antibody

别 名: p65; NFKB3

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: RELA (Phospho-Thr505)

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processed such as inflammation, in mmunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct p



| | references for different kappa-B sites that they can bind with distinguishable affinity and specificity. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-k appa-B inhibitor (I-kappa-B) family. In a conventiona I activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different acti vators, subsequently degraded thus liberating the ac tive NF-kappa-B heterodimeric p65-p50 and p65-c-R el complexes are transcriptional activators. The inhibitory effect of I-kappa-B upon NF-kappa-B the cytopl asm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-kappa-B complex. |
|--------------------------|--|
| Applications: | WB, IHC |
| Name of antibody: | RELA (Phospho-Thr505) |
| Immunogen: | Synthetic peptide of human RELA (Phospho-Thr505) |
| Full name: | v-rel reticuloendotheliosis viral oncogene homolog A (avian) (Phospho-Thr505) |
| Synonyms: | p65; NFKB3 |
| SwissProt: | Q04206 |
| IHC positive control: | Human breast carcinoma |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 65 kDa |
| WB Positive control: | HL60 cells treated with TNF- α |
| WB Recommended dilution: | 500-1000 |





