

兔抗 NFATC4 多克隆抗体

中文名称: 兔抗 NFATC4 多克隆抗体

英文名称: Anti-NFATC4 rabbit polyclonal antibody

别 名: NFAT3; NF-ATc4

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: NFATC4

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

The product of this gene is a member of the nuclear fact ors of activated T cells DNA-binding transcription comple x. This complex consists of at least two components: a pr eexisting cytosolic component that translocates to the nu cleus upon T cell receptor (TCR) stimulation and an induc ible nuclear component. Other members of this family of nuclear factors of activated T cells also participate in the formation of this complex. The product of this gene plays a role in the inducible expression of cytokine genes in T cells, especially in the induction of the IL-2 and IL-4. Alte rnatively spliced transcript variants encoding different isof



| | orms have been noted for this gene. |
|------------------------------------|--|
| Applications: | ELISA, WB |
| Name of antibody: | NFATC4 |
| Immunogen: | Synthetic peptide of human NFATC4 |
| Full name: | Nuclear factor of activated T-cells, cytoplasmic, calcineurin -dependent 4 |
| Synonyms : | NFAT3; NF-ATc4 |
| SwissProt: | Q14934 |
| ELISA Recommended dilution: | 2000-5000 |
| WB Predicted band size: | 102 kDa |
| WB Positive control: | NIH/3T3 and Jurkat cells |
| WB Recommended dilution: | 500-2000 |

