

兔抗 UBE2D3 多克隆抗体

- 中文名称：兔抗 UBE2D3 多克隆抗体
- 英文名称：Anti-UBE2D3 rabbit polyclonal antibody
- 别名：UBC4/5; UBCH5C; E2(17)KB3
- 储存：冷冻 (-20℃)
- 宿主：Rabbit
- 抗原：UBE2D3
- 反应种属：Human, mouse, Rat
- 相关类别：一抗
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme functions in the ubiquitination of the tumor-suppressor protein p53, which is induced by an E3 ubiquitin-protein ligase. Multiple spliced transcript variants have been found

	for this gene, but the full-length nature of some variants has not been determined.
Applications:	ELISA, WB, IHC
Name of antibody:	UBE2D3
Immunogen:	Full length fusion protein
Full name:	ubiquitin conjugating enzyme E2D 3
Synonyms:	UBC4/5; UBCH5C; E2(17)KB3
SwissProt:	P61077
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer and human tonsil
IHC Recommend dilution:	25-100
WB Predicted band size:	17 kDa
WB Positive control:	NIH/3T3 cell lysate
WB Recommended dilution:	200-1000





