

兔抗 NKD2 多克隆抗体

中文名称：兔抗 NKD2 多克隆抗体

英文名称：Anti-NKD2 rabbit polyclonal antibody

别名：naked cuticle homolog 2; Naked2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：NKD2

反应种属：Human, Mouse


标记物：Unconjugate



克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | This gene encodes a member of a family of proteins that function as negative regulators of Wnt receptor signaling through interaction with Dishevelled family members. The encoded protein participates in the delivery of transforming growth factor alpha-containing vesicles to the cell membrane. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | NKD2 |
| Immunogen: | Fusion protein of human NKD2 |
| Full name: | naked cuticle homolog 2 |
| Synonyms: | Naked2 |
| SwissProt: | Q969F2 |

| | |
|------------------------------------|--|
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human tonsil and Human prostate cancer |
| IHC Recommend dilution: | 40-200 |
| WB Predicted band size: | 50 kDa |
| WB Positive control: | Mouse Pancreas tissue lysate |
| WB Recommended dilution: | 500-2000 |





mlbio 梅联生物
Good elisakit producers