

## 兔抗 UBR7 多克隆抗体

中文名称：兔抗 UBR7 多克隆抗体

英文名称：Anti-UBR7 rabbit polyclonal antibody

别名：ubiquitin protein ligase E3 component n-recognin 7 (putative); C14orf130

相关类别：一抗

抗原：UBR7

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

#### Background:

Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). Ubr7 (ubiquitin protein ligase E3 component n-recognin 7), also known as C14orf130 or N-recognin-7, is a 425 amino acid protein that contains one UBR-type zinc finger and one PHD zinc finger. Participating in protein modification events within the N-end rule p

	athway, Ubr7 functions as an E3 ubiquitin-protein ligase that recognizes and binds proteins that contain destabilizing N-terminal residues, thereby leading to their ubiquitination and subsequent degradation.
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	UBR7
<b>Immunogen:</b>	Fusion protein of human UBR7
<b>Full name:</b>	ubiquitin protein ligase E3 component n-recogin 7 (putative)
<b>Synonyms:</b>	C14orf130
<b>SwissProt:</b>	Q8N806
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human colorectal cancer;Human liver cancer
<b>IHC Recommend dilution:</b>	30-150
<b>WB Predicted band size:</b>	48 kDa
<b>WB Positive control:</b>	Hela cell lysate
<b>WB Recommended dilution:</b>	200-1000





