

兔抗 UCHL5 多克隆抗体

中文名称：兔抗 UCHL5 多克隆抗体

英文名称： Anti-UCHL5 rabbit polyclonal antibody

别名： UCH37; CGI-70; INO80R; UCH-L5

相关类别： 一抗

抗原： UCHL5

储存： 冷冻（-20℃）

宿主： Rabbit

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | Ubiquitin carboxyl-terminal hydrolase isozyme L5 is an enzyme that in humans is encoded by the UCHL5 gene. Protease that specifically cleaves 'Lys-48'-linked polyubiquitin chains. Deubiquitinating enzyme associated with the 19S regulatory subunit of the 26S proteasome. Putative regulatory component of the INO80 complex; however is inactive in the INO80 complex and is activated by a transient interaction of the INO80 complex with the proteasome via ADRM1. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | UCHL5 |
| Immunogen: | Fusion protein of human UCHL5 |
| Full name: | ubiquitin carboxyl-terminal hydrolase L5 |

| | |
|------------------------------------|--|
| Synonyms : | UCH37; CGI-70; INO80R; UCH-L5 |
| SwissProt: | Q9Y5K5 |
| ELISA Recommended dilution: | 1000-2000 |
| IHC positive control: | Human thyroid cancer and human cervical cancer |
| IHC Recommend dilution: | 25-100 |
| WB Predicted band size: | 38 kDa |
| WB Positive control: | Human fetal kidney tissue and 293T cells |
| WB Recommended dilution: | 200-1000 |



