

兔抗 RPA2(Ab-21) 多克隆抗体

中文名称：兔抗 RPA2(Ab-21) 多克隆抗体

英文名称： Anti-RPA2(Ab-21) rabbit polyclonal antibody

别名： replication protein A2, 32kDa

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： RPA2(Ab-21)

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing.

ling and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage the RPA complex controls DNA repair and DNA damage checkpoint activation. It is required for the recruitment of the DNA double-strand break repair factors RAD52 and RAD51 to chromatin in response to DNA damage. Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair. Plays also a role in base excision repair (BER) probably through interaction with UNG. Through RPA3 may activate CHEK1 and play a role in replicati

	<p>on checkpoint control. Also recruits SMA RCAL1/HARP which is involved in replicati on fork restart to sites of DNA damage. May also play a role in telomere mainten ance.</p>
Applications:	WB, IHC
Name of antibody:	RPA2(Ab-21)
Immunogen:	Synthesized non-phosphopeptide derived from human RFA2 around the phosphoryl ation site of threonine 21 (G-Y-T(p)-Q-S).
Full name:	replication protein A2, 32kDa
Synonyms :	replication protein A2, 32kDa
SwissProt:	P15927
IHC positive control:	Human lung carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	29 kDa
WB Positive control:	K562 cells lysate
WB Recommended dilution:	500-3000

