

兔抗 RPS6KA5(Ab-212) 多克隆抗体

中文名称：兔抗 RPS6KA5(Ab-212) 多克隆抗体

英文名称： Anti-RPS6KA5(Ab-212) rabbit polyclonal antibody

别名： MSK1; RLPK; MSPK1

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： RPS6KA5(Ab-212)

反应种属： Human Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

Serine/threonine-protein kinase that is required for the mitogen or stress-induced phosphorylation of the transcription factors CREB1 and ATF1 and for the regulation of the transcription factors REL

A, STAT3 and ETV1/ER81, and that contributes to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes. Phosphorylates CREB1 and ATF1 in response to mitogenic or stress stimuli such as UV-C irradiation, epidermal growth factor (EGF) and anisomycin. Plays an essential role in the control of RELA transcriptional activity in response to TNF and upon glucocorticoid, associates in the cytoplasm with the glucocorticoid receptor NR3C1 and contributes to RELA inhibition and repression of inflammatory gene expression. In skeletal myoblasts is required for phosphorylation of RELA at 'Ser-276' during oxidative stress. In erythropoietin-stimulated cells, is necessary for the 'Ser-727' phosphorylation of STAT3 and reg

ulation of its transcriptional potential. P
hosphorylates ETV1/ER81 at 'Ser-191' an
d 'Ser-216', and thereby regulates its ab
ility to stimulate transcription, which ma
y be important during development and
breast tumor formation. Directly repress
es transcription via phosphorylation of '
Ser-1' of histone H2A. Phosphorylates 'S
er-10' of histone H3 in response to mit
ogenics, stress stimuli and EGF, which r
esults in the transcriptional activation of
several immediate early genes, including
proto-oncogenes c-fos/FOS and c-jun/J
UN. May also phosphorylate 'Ser-28' of
histone H3. Mediates the mitogen- and
stress-induced phosphorylation of high
mobility group protein 1 (HMGN1/HMG
14). In lipopolysaccharide-stimulated pri
mary macrophages, acts downstream of

	<p>the Toll-like receptor TLR4 to limit the production of pro-inflammatory cytokines. Functions probably by inducing transcription of the MAP kinase phosphatase DUSP1 and the anti-inflammatory cytokine interleukin 10 (IL10), via CREB1 and ATF1 transcription factors. Plays a role in neuronal cell death by mediating the downstream effects of excitotoxic injury.</p>
Applications:	WB, IHC
Name of antibody:	RPS6KA5(Ab-212)
Immunogen:	Synthesized peptide derived from Internal of human MSK1.
Full name:	ribosomal protein S6 kinase, 90kDa, polypeptide 5
Synonyms :	MSK1; RLPK; MSPK1
SwissProt:	O75582
IHC positive control:	Human colon carcinoma tissue
IHC Recommend dilution:	50-100

WB Predicted band size:	67 kDa
WB Positive control:	Mouse brain tissue lysate
WB Recommended dilution:	500-3000



