

## 兔抗 NXF3 多克隆抗体

中文名称: 兔抗 NXF3 多克隆抗体

英文名称: Anti-NXF3 rabbit polyclonal antibody

抗 原: NXF3

储 存: 冷冻 (-20℃) 避光

宿 主: Rabbit

反应种属: Human

相关类别: 一抗

标 记 物: Unconjugate

克隆类型: Unconjugate

技术规格

|                          |   |
|--------------------------|---|
| <b>Background:</b>       | This gene is one member of a family of nuclear RNA export factor genes. Common domain features of this family are a noncanonical RNP-type RNA-binding domain (RBD), 4 leucine-rich repeats (LRRs), a nuclear transport factor 2 (NTF2)-like domain that allows heterodimerization with NTF2-related export protein-1 (NXT1), and a ubiquitin-associated domain that mediates interactions with nucleoporins. The LRRs and NTF2-like domains are required for export activity. Alternative splicing seems to be a common mechanism in this gene family. The encoded protein of this gene has shortened LRR and ubiquitin-associated domains and its RBD is unable to bind RNA. It is located in the nucleoplasm but is not associated with either the nuclear envelope or the nucleolus. |
| <b>Applications:</b>     | WB  |
| <b>Name of antibody:</b> | NXF3  |

|                                 |  |
|---------------------------------|--|
| <b>Immunogen:</b>               | Synthesized peptide derived from internal of human NXF3. |
| <b>Full name:</b>               | nuclear RNA export factor 3                              |
| <b>Synonyms :</b>               | 0  |
| <b>SwissProt:</b>               | Q9H4D5   |
| <b>WB Predicted band size:</b>  | 60 kDa   |
| <b>WB Positive control:</b>     | MCF-7 cells lysate                                       |
| <b>WB Recommended dilution:</b> | 500-3000   |

