

兔抗 ITCH(Phospho-Tyr420) 多克隆抗体

- 中文名称： 兔抗 ITCH(Phospho-Tyr420) 多克隆抗体
- 英文名称： Anti-ITCH(Phospho-Tyr420) rabbit polyclonal antibody
- 别名： AIF4; AIP4; ADMFD; NAPP1; dJ468O1.1
- 相关类别： 一抗
- 储存： 冷冻（-20℃） 避光
- 宿主： Rabbit
- 抗原： ITCH(Phospho-Tyr420)
- 反应种属： Human Mouse
- 标记物： Unconjugate
- 克隆类型： Unconjugate

技术规格

Background:	This gene encodes a member of the Nedd4 family of HECT domain E3 ubiquitin ligases. HECT domain E3 ubiquitin ligases transfer ubiquitin from E2 ubiquitin-conjugating enzymes to protein substrates, thus targeting specific proteins for lysosomal degradation. The encoded protein plays a role in multiple cellular processes including erythroid and lymphoid cell differentiation and the regulation of immune responses. Mutations in this gene are a cause of syndromic multisystem autoimmune disease. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Mar 2012]
Applications:	IHC
Name of antibody:	ITCH(Phospho-Tyr420)

Immunogen:	Peptide sequence around phosphorylation site of tyrosine 420(F-I-Y(p)-G-N) derived from Human ITCH .
Full name:	itchy E3 ubiquitin protein ligase
Synonyms :	AIF4; AIP4; ADMFD; NAPP1; dJ468O1.1
SwissProt:	Q96J02
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100

