

兔抗 SDC1 多克隆抗体

中文名称：兔抗 SDC1 多克隆抗体

英文名称：Anti-SDC1 rabbit polyclonal antibody

别名：SDC; CD138; SYND1; syndecan

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：SDC1

反应种属：Human

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:

The protein encoded by this gene is a transmembrane (type I) heparan sulfate proteoglycan and is a member of the syndecan proteoglycan family. The syndecans mediate cell binding, cell signaling, and cytoskeletal organization and syndecan receptors are required for internalization of the HIV-1 tat protein. The syndecan-1 protein functions as an integral membrane protein and participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. Altered syndecan-1 expression has been detected in several different

	tumor types. While several transcript variants may exist for this gene, the full-length nature of only two have been described to date. These two represent the major variants of this gene and encode the same protein.
Applications:	WB
Name of antibody:	SDC1
Immunogen:	Synthesized peptide derived from internal of human SDC1.
Full name:	syndecan 1
Synonyms :	SDC; CD138; SYND1; syndecan
SwissProt:	P18827
WB Predicted band size:	32 kDa
WB Positive control:	Hela cells lysate
WB Recommended dilution:	500-3000

