

兔抗 SDC2 多克隆抗体

- 中文名称: 兔抗 SDC2 多克隆抗体
- 英文名称: Anti-SDC2 rabbit polyclonal antibody
- 别名: HSPG; CD362; HSPG1; SYND2
- 相关类别: 一抗
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 抗原: SDC2
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

| | The protein encoded by this gene is a transme |
|-------------|--|
| | mbrane (type I) heparan sulfate proteoglycan an |
| | d is a member of the syndecan proteoglycan fa |
| | mily. The syndecans mediate cell binding, cell si |
| | gnaling, and cytoskeletal organization and synde |
| Background: | can receptors are required for internalization of |
| | the HIV-1 tat protein. The syndecan-2 protein fu |
| | nctions as an integral membrane protein and pa |
| | rticipates in cell proliferation, cell migration and |
| | cell-matrix interactions via its receptor for extrac |
| | ellular matrix proteins. Altered syndecan-2 expre |



| | ssion has been detected in several different tum or types. |
|-----------------------------|--|
| Applications: | ELISA, IHC |
| Name of antibody: | SDC2 |
| Immunogen: | Synthetic peptide of human SDC2 |
| Full name: | syndecan 2 |
| Synonyms: | HSPG; CD362; HSPG1; SYND2 |
| SwissProt: | P34741 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human colorectal cancer and human brain |
| IHC Recommend dilution: | 25-100 |





