

## 兔抗 WASF2 多克隆抗体

中文名称: 兔抗 WASF2 多克隆抗体

英文名称: Anti-WASF2 rabbit polyclonal antibody

别 名: IMD2; SCAR2; WASF4; WAVE2; dJ393P12.2

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: WASF2

反应种属: Human, Mouse

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

This gene encodes a member of the Wiskott-Aldr ich syndrome protein family. The gene product is a protein that forms a multiprotein complex that links receptor kinases and actin. Binding to actin occurs through a C-terminal verprolin homology domain in all family members. The multiprotein c omplex serves to tranduce signals that involve ch anges in cell shape, motility or function. The pub lished map location (PMID:10381382) has been ch anged based on recent genomic sequence comparisons, which indicate that the expressed gene is



	located on chromosome 1, and a pseudogene m ay be located on chromosome X.
Applications:	ELISA, WB, IHC
Name of antibody:	WASF2
Immunogen:	Synthetic peptide of human WASF2
Full name:	WAS protein family, member 2
Synonyms:	IMD2; SCAR2; WASF4; WAVE2; dJ393P12.2
SwissProt:	Q9Y6W5
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human brain and Human esophagus cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	54 kDa
WB Positive control:	Mouse lung tissue
WB Recommended dilution:	200-1000





