

兔抗 SFXN1 多克隆抗体

- 中文名称: 兔抗 SFXN1 多克隆抗体
- 英文名称: Anti-SFXN1 rabbit polyclonal antibody
- 抗原: SFXN1
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 相关类别: 一抗
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

Background:	The sideroflexin (SFXN) family is comprised of SF XN1, SFXN2, SFXN3, SFXN4 and SFXN5. SFXN1, a lso designated tricarboxylate carrier protein TCC, is the most highly characterized family member. The ubiquitously expressed SFXN1 protein resides as an integral protein of the mitochondrial inner membrane. It functions as an essential compone nt of the shuttle system that transports mitocho ndrial acetyl-CoA into the cytosol, where lipogen esis occurs. The SFXN1 gene is mutated in flexed -tail (f/f) mice, which display axial skeletal abnor malities and a transient embryonic and neonatal anemia characterized by pathologic intramitochon
-------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



	diral iron deposits in erythrocytes. Therefore, SFX N1 is also thought to facilitate the transport of a component required for iron utilization into mi tochondria. All SFXN family members show expre ssion in pancreatic islet cells. SFXN5 displays a ci trate transport activity and is primarily expressed in brain.
Applications:	ELISA, WB
Name of antibody:	SFXN1
Immunogen:	Fusion protein of human SFXN1
Full name:	sideroflexin 1
SwissProt:	Q9H9B4
ELISA Recommended dilution:	5000-10000
WB Predicted band size:	36 kDa
WB Positive control:	Human fetal liver tissue and Human cerebrum tis sue
WB Recommended dilution:	200-1000





