

兔抗 SIGLEC12 多克隆抗体

中文名称：兔抗 SIGLEC12 多克隆抗体

英文名称： Anti-SIGLEC12 rabbit ployclonal antibody

别名： S2V; SLG; SIGLECL1; Siglec-XII

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： SIGLEC12

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

Sialic acid-binding immunoglobulin-like lectins (SIGLECs) are a family of cell surface proteins belonging to the immunoglobulin superfamily. They mediate protein-carbohydrate interactions by selectively binding to different sialic acid moieties present on glycolipids and glycoproteins. This gene encodes a member of the SIGLEC3-like subfamily of SIGLECs. Members of this subfamily are characterized by an extracellular V-set immunoglobulin-like domain followed by two C2-set immunoglobulin-like domains, and the cytoplasmic tyrosine-based motifs ITIM an

	d SLAM-like.
Applications:	ELISA, WB, IHC
Name of antibody:	SIGLEC12
Immunogen:	Fusion protein of human SIGLEC12
Full name:	sialic acid binding Ig-like lectin 12 (gene/pseudogene)
Synonyms :	S2V; SLG; SIGLECL1; Siglec-XII
SwissProt:	Q96PQ1
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human prostate cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	65 kDa
WB Positive control:	Mouse bladder tissue
WB Recommended dilution:	200-1000



