

兔抗 KCNK18 多克隆抗体

中文名称：兔抗 KCNK18 多克隆抗体

英文名称：Anti-KCNK18 rabbit polyclonal antibody

别名：potassium two pore domain channel subfamily K member 18; TRIK; MGR13; TRESK; TRESK2; K2p18.1; TRESK-2

相关类别：一抗

储存：冷冻 (-20℃)

宿主：Rabbit

抗原：KCNK18

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Potassium channels play a role in many cellular processes including maintenance of the action potential, muscle contraction, hormone secretion, osmotic regulation, and ion flow. This gene encodes a member of the superfamily of potassium channel proteins containing two pore-forming P domains and the encoded protein functions as an outward rectifying potassium channel. A mutation in this gene has been found to be associated with migraine with aura.
Applications:	ELISA, WB, IHC

Name of antibody:	KCNK18
Immunogen:	Synthetic peptide of human KCNK18
Full name:	potassium two pore domain channel subfamily K member 18
Synonyms:	TRIK; MGR13; TRESK; TRESK2; K2p18.1; TRESK-2
SwissProt:	Q7Z418
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	70-350
WB Predicted band size:	44 kDa
WB Positive control:	TM4 cell lysate
WB Recommended dilution:	500-2000



