

兔抗 ZAP70 多克隆抗体

中文名称：兔抗 ZAP70 多克隆抗体

英文名称：Anti-ZAP70 rabbit polyclonal antibody

别名：zeta chain of T cell receptor associated protein kinase 70; SRK; STD; TZK; STCD; IMD48; ADMIO2; ZAP-70

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：ZAP70

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe

	combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	ZAP70
Immunogen:	Fusion protein of human ZAP70
Full name:	zeta chain of T cell receptor associated protein kinase 70
Synonyms:	SRK; STD; TZK; STCD; IMD48; ADMIO2; ZAP-70
SwissProt:	P43403
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human cervical cancer
IHC Recommend dilution:	50-300
WB Predicted band size:	70 kDa
WB Positive control:	Mouse liver tissue and Jurkat cell lysates
WB Recommended dilution:	500-2000



