

## 兔抗 PROC 多克隆抗体

中文名称：兔抗 PROC 多克隆抗体

英文名称：Anti-PROC rabbit polyclonal antibody

别名：protein C, inactivator of coagulation factors Va and VIIIa; PC; APC; PROC1; THPH3; THPH4

相关类别：一抗

抗原：PROC

储存：冷冻 (-20℃)

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

**Background:**

This gene encodes a vitamin K-dependent plasma glycoprotein. The encoded protein is cleaved to its activated form by the thrombin-thrombomodulin complex. This activated form contains a serine protease domain and functions in degradation of the activated forms of coagulation factors V and VIII. Mutations in this gene have been associated with thrombophilia due to protein C deficiency, neonatal purpura fulminans, and recurrent venous thrombosis.

<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	PROC
<b>Immunogen:</b>	Synthetic peptide of human PROC
<b>Full name:</b>	protein C, inactivator of coagulation factors Va and VIIIa
<b>Synonyms:</b>	PC; APC; PROC1; THPH3; THPH4
<b>SwissProt:</b>	P04070
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human prostate cancer and Human gastric cancer
<b>IHC Recommend dilution:</b>	50-100
<b>WB Predicted band size:</b>	52 kDa
<b>WB Positive control:</b>	SKOV3 cell lysate
<b>WB Recommended dilution:</b>	500-2000



