

兔抗 L3MBTL2 多克隆抗体

中文名称：兔抗 L3MBTL2 多克隆抗体

英文名称：Anti-L3MBTL2 rabbit polyclonal antibody

别名：L3MBTL2, polycomb repressive complex 1 subunit; L3MBT; H-I(3)mbt-l

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：L3MBTL2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Putative Polycomb group (PcG) protein. PcG proteins maintain the transcriptionally repressive state of genes, probably via a modification of chromatin, rendering it heritably changed in its expressibility. Its association with a chromatin-remodeling complex suggests that it may contribute to prevent expression of genes that trigger the cell into mitosis. Binds to monomethylated and dimethylated 'Lys-20' on histone H4. Binds histone H3 peptides that are monomethylated or dimethylated on 'Lys-4', 'Lys-9' or 'Lys-27'.
Applications:	ELISA, WB, IHC

Name of antibody:	L3MBTL2
Immunogen:	Fusion protein of human L3MBTL2
Full name:	L3MBTL2, polycomb repressive complex 1 subunit
Synonyms:	L3MBT; H-I(3)mbt-l
SwissProt:	Q969R5
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human thyroid cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	79 kDa
WB Positive control:	Mouse thymus tissue lysate
WB Recommended dilution:	500-2000



