

## 小鼠抗 CCND2 单克隆抗体

中文名称: 小鼠抗 CCND2 单克隆抗体

英文名称: Anti-CCND2 mouse monoclonal antibody

别 名: cyclin D2; MPPH3; KIAK0002

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Mouse

抗 原: CCND2

反应种属: Human

标 记 物: Unconjugate

克隆类型: Mouse Monoclonal

## 技术规格

**Background:** 

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterize d by a dramatic periodicity in protein abundance throug h the cell cycle. Cyclins function as regulators of CDK ki nases. Different cyclins exhibit distinct expression and de gradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a comp lex with CDK4 or CDK6 and functions as a regulatory su bunit of the complex, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tu



	mor suppressor protein Rb. Knockout studies of the hom ologous gene in mouse suggest the essential roles of thi s gene in ovarian granulosa and germ cell proliferation. High level expression of this gene was observed in ovari an and testicular tumors. Mutations in this gene are ass ociated with megalencephaly-polymicrogyria-polydactyly-h ydrocephalus syndrome 3 (MPPH3).
Applications:	WB
Name of antibody:	CCND2
Immunogen:	Fusion protein of human CCND2
Full name:	cyclin D2
Synonyms:	МРРН3; КІАК0002
SwissProt:	P30279
WB Predicted band size:	33 kDa
WB Positive control:	CHO-K1 cell and CHO-K1 cell transfected by Cyclin D2-fr agment EGFP fusion protein lysates
WB Recommended dilution:	1000-5000

