

兔抗 STAT5A (Phospho-Tyr694) 多克隆抗体

中文名称: 兔抗 STAT5A (Phospho-Tyr694) 多克隆抗体

英文名称: Anti-STAT5A (Phospho-Tyr694) rabbit polyclonal antibody

别 名: MGF; STAT5

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: STAT5A (Phospho-Tyr694)

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or he terodimers that translocate to the cell nucleus where they a ct as transcription activators. This protein is activated by, an d mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and differe nt growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is in dependent of cell stimulus and has been shown to be esse



	ntial for the tumorigenesis. The mouse counterpart of this g ene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in ce lls.
Applications:	WB, IHC
Name of antibody:	STAT5A (Phospho-Tyr694)
Immunogen:	Synthetic peptide of human STAT5A (Phospho-Tyr694)
Full name:	signal transducer and activator of transcription 5A (Phospho -Tyr694)
Synonyms:	MGF; STAT5
SwissProt:	P42229
IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	91 kDa
WB Positive control:	K562 cells treated with G-CSF
WB Recommended dilution:	500-1000





