

## 小鼠抗 DDB1 单克隆抗体

中文名称： 小鼠抗 DDB1 单克隆抗体

英文名称： Anti-DDB1 mouse monoclonal antibody

别名： damage specific DNA binding protein 1; XPE; DDBA; XAP1; XPCE; XPE-BF; UV-DDB1

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Mouse

抗原： DDB1

反应种属： Human, Mouse, Rat, Monkey

标记物： Unconjugate

克隆类型： Mouse Monoclonal

### 技术规格

#### Background:

The protein encoded by this gene is the large subunit (p127) of the heterodimeric DNA damage-binding (DDB) complex while another protein (p48) forms the small subunit. This protein complex functions in nucleotide-excision repair and binds to DNA following UV damage. Defective activity of this complex causes the repair defect in patients with xeroderma pigmentosum complementation group E (XPE) - an autosomal recessive disorder characterized by photosensitivity and early onset of carcinomas. However, it remains for mutation analysis to demonstrate whether the defect in XPE patients is in this gene or the gene encoding the small s

	ubunit. In addition, Best vitelliform muscular dystrophy is mapped to the same region as this gene on 11q, but no sequence alternations of this gene are demonstrated in Best disease patients. The protein encoded by this gene also functions as an adaptor molecule for the cullin 4 (CUL4) ubiquitin E3 ligase complex by facilitating the binding of substrates to this complex and the ubiquitination of proteins.
<b>Applications:</b>	WB
<b>Name of antibody:</b>	DDB1
<b>Immunogen:</b>	Fusion protein of human DDB1
<b>Full name:</b>	damage specific DNA binding protein 1
<b>Synonyms:</b>	XPE; DDBA; XAP1; XPCE; XPE-BF; UV-DDB1
<b>SwissProt:</b>	Q16531
<b>WB Predicted band size:</b>	127 kDa
<b>WB Positive control:</b>	MCF7, C6, 3T3, COS7 and Hela cell lysates
<b>WB Recommended dilution:</b>	1000-5000

