

小鼠抗 FCGR3A 单克隆抗体

- 中文名称: 小鼠抗 FCGR3A 单克隆抗体
- 英文名称: Anti-FCGR3A mouse monoclonal antibody

别 名: Fc fragment of IgG receptor IIIa; CD16; FCG3; CD16A; FCGR3; IGFR3; IMD20; FCR-10; FCRIII; FCGRIII; FCRIIIA

- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 宿 主: Mouse
- 抗 原: FCGR3A
- 反应种属: Mouse
- 标记物: Unconjugate
- 克隆类型: mouse monoclonal

技术规格

Background:	This gene encodes a receptor for the Fc portion of imm unoglobulin G, and it is involved in the removal of anti gen-antibody complexes from the circulation, as well as other other antibody-dependent responses. This gene (F CGR3A) is highly similar to another nearby gene (FCGR3 B) located on chromosome 1. The receptor encoded by this gene is expressed on natural killer (NK) cells as an i ntegral membrane glycoprotein anchored through a tran smembrane peptide, whereas FCGR3B is expressed on p olymorphonuclear neutrophils (PMN) where the receptor
-------------	--



	 is anchored through a phosphatidylinositol (PI) linkage. Mutations in this gene have been linked to susceptibility to recurrent viral infections, susceptibility to systemic lup us erythematosus, and alloimmune neonatal neutropenia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Applications:	WB
Name of antibody:	FCGR3A
Immunogen:	Fusion protein of human FCGR3A
Full name:	Fc fragment of IgG receptor IIIa
Synonyms:	CD16; FCG3; CD16A; FCGR3; IGFR3; IMD20; FCR-10; FCRII I; FCGRIII; FCRIIIA
SwissProt:	P08637
WB Predicted band size:	29 kDa
WB Positive control:	Mouse liver tissue lysate
WB Recommended dilution:	1000-5000

