

兔抗 TBC1D4 多克隆抗体

中文名称：兔抗 TBC1D4 多克隆抗体

英文名称：Anti-TBC1D4 rabbit polyclonal antibody

别名：AS160

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：TBC1D4

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

TBC1 domain family member 4 (TBC1D4), also designated AS160, can be insulin- and/or AKT1-induced. Insulin-stimulated phosphorylation is required for GLUT4 translocation. TBC1D4 may play a role as a GTPase activating protein for proteins in the Rab family. It is expressed primarily in skeletal muscle and heart, as well as spleen, lymph node and leukocytes. Defects in the TBC1D4 gene may cause atopic dermatitis (AD), sometimes referred to as eczema, an atopic chronic skin disease. The skin of affected individuals reacts to irritants or allergens and beco

	mes red, flaky and itchy. The skin is also more vulnerable to inflammations, and symptoms can grow or disappear over time.
Applications:	ELISA, IHC
Name of antibody:	TBC1D4
Immunogen:	Synthetic peptide of human TBC1D4
Full name:	TBC1 domain family, member 4
Synonyms :	AS160
SwissProt:	O60343
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human liver cancer and Human brain
IHC Recommend dilution:	25-100

