

兔抗 LZTFL1 多克隆抗体

- 中文名称：兔抗 LZTFL1 多克隆抗体
- 英文名称：Anti-LZTFL1 rabbit polyclonal antibody
- 别名：BBS17
- 相关类别：一抗
- 抗原：LZTFL1
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes a ubiquitously expressed protein that localizes to the cytoplasm. This protein interacts with Bardet-Biedl Syndrome (BBS) proteins and, through its interaction with BBS protein complexes, regulates protein trafficking to the ciliary membrane. Nonsense mutations in this gene cause a form of Bardet-Biedl Syndrome; a ciliopathy characterized in part by polydactyly, obesity, cognitive impairment, hypogonadism, and kidney failure. This gene may also function as a tumor suppressor; possibly by interacting with E-cadherin and the actin cytoskeleton and thereby regulating the transition of epithelial cell

	s to mesenchymal cells. Alternative splicing of this gene results in multiple transcript variants.
Applications:	ELISA, WB, IHC
Name of antibody:	LZTFL1
Immunogen:	Fusion protein of human LZTFL1
Full name:	leucine zipper transcription factor-like 1
Synonyms:	BBS17
SwissProt:	Q9NQ48
IHC positive control:	Human gastric cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	35 kDa
WB Positive control:	Human transitional cells carcinoma tissue, 293T and A172 cells, human testis tissue and Hela cells
WB Recommended dilution:	500-2000



