

小鼠抗 GORAB 单克隆抗体

| | |
|-------|--------------------------------------|
| 中文名称: | 小鼠抗 GORAB 单克隆抗体 |
| 英文名称: | Anti-GORAB mouse monoclonal antibody |
| 别名: | GO; NTKLBP1; SCYL1BP1 |
| 抗原 | GORAB |
| 储存 | 冷冻 (-20℃) 避光 |
| 宿主 | Mouse |
| 反应种属 | Human , Dog , Rat , Monkey , Mouse |
| 标记物 | Unconjugate |
| 相关类别 | 一抗 |
| 克隆类型 | mouse monoclonal |

技术规格

| | |
|----------------------|--|
| Background: | This gene encodes a member of the golgin family, a group of coiled-coil proteins localized to the Golgi. The encoded protein may function in the secretory pathway. The encoded protein, which also localizes to the cytoplasm, was identified by interactions with the N-terminal kinase-like protein, and thus it may function in mitosis. Mutations in this gene have been associated with geroderma osteodysplastica. Alternatively spliced transcript variants have been described. [provided by RefSeq]. |
| Applications: | WB, IHC |

| | |
|---------------------------------|--|
| Name of antibody: | GORAB |
| Immunogen: | Fusion protein of human GORAB |
| Full name: | golgin, RAB6-interacting (GORAB), transcript variant 1 |
| Synonyms: | GO; NTKLBP1; SCYL1BP1 |
| SwissProt: | Q5T7V8 |
| IHC positive control: | human pancreas tissue and adenocarcinoma of human endometrium tissue; carcinoma of human liver tissue and carcinoma of human prostate tissue |
| IHC Recommend dilution: | 30-150 |
| WB Predicted band size: | 45 kDa |
| WB Positive control: | HepG2, Hela, SVT2, A549, COS7, Jurkat, MDCK, PC12, MCF-7 cell lysates |
| WB Recommended dilution: | 200-1000 |