

## 小鼠抗 LAMC1 单克隆抗体

中文名称： 小鼠抗 LAMC1 单克隆抗体

英文名称： Anti-LAMC1 mouse monoclonal antibody

相关类别： 一抗

储 存： 冷冻（-20℃）

宿 主： Mouse

抗 原： LAMC1

反应种属： Human

标 记 物： Unconjugate

克隆类型： mouse monoclonal

### 技术规格

**Background:**

Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminin is composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively), have a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. Several isoforms of each chain have been described. Different alpha, beta and gamma chain isoforms combine to give rise to different heterotrimeric lamin

	<p>in isoforms which are designated by Arabic numerals in the order of their discovery, i.e. alpha1beta1gamma1 heterotrimer is laminin 1. The biological functions of the different chains and trimer molecules are largely unknown, but some of the chains have been shown to differ with respect to their tissue distribution, presumably reflecting diverse functions in vivo. This gene encodes the gamma chain isoform laminin, gamma 1. The gamma 1 chain, formerly thought to be a beta chain, contains structural domains similar to beta chains, however, lacks the short alpha region separating domains I and II. The structural organization of this gene also suggested that it had diverged considerably from the beta chain genes. Embryos of transgenic mice in which both alleles of the gamma 1 chain gene were inactivated by homologous recombination, lacked basement membranes, indicating that laminin, gamma 1 chain is necessary for laminin heterotrimer assembly. It has been inferred by analogy with the strikingly similar 3' UTR sequence in mouse laminin gamma 1 cDNA, that multiple polyadenylation sites are utilized in human to generate the 2 different sized mRNAs (5.5 and 7.5 kb) seen on Northern analysis.</p>
<b>Applications:</b>	ELISA, WB
<b>Name of antibody:</b>	LAMC1
<b>Immunogen:</b>	Fusion protein of human LAMC1
<b>Full name:</b>	laminin subunit gamma 1
<b>Synonyms:</b>	LAMB2; Laminin B2 chain
<b>SwissProt:</b>	P11047
<b>WB Predicted band size:</b>	178 kDa
<b>WB Positive control:</b>	SW480 whole cell lysate
<b>WB Recommended dilution:</b>	500-2000

