

小鼠抗 PSMC2 单克隆抗体

中文名称： 小鼠抗 PSMC2 单克隆抗体

英文名称： Anti-PSMC2 mouse monoclonal antibody

相关类别： 一抗

储 存： 冷冻（-20℃）

宿 主： Mouse

抗 原： PSMC2

反应种属： Human

标 记 物： Unconjugate

克隆类型： mouse monoclonal

技术规格

Background:

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a

	member of the triple-A family of ATPases which have a chaperone-like activity. This subunit has been shown to interact with several of the basal transcription factors so, in addition to participation in proteasome functions, this subunit may participate in the regulation of transcription. This subunit may also compete with PSMC3 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. Alternative splicing results in multiple transcript variants encoding distinct isoforms.
Applications:	ELISA, WB
Name of antibody:	PSMC2
Immunogen:	Fusion protein of human PSMC2
Full name:	proteasome 26S subunit, ATPase 2
Synonyms:	S7; MSS1; Nbla10058
SwissProt:	P35998
WB Predicted band size:	49 kDa
WB Positive control:	HeLa whole cell lysate
WB Recommended dilution:	1000-5000

