

兔抗 ZFAND5 多克隆抗体

中文名称: 兔抗 ZFAND5 多克隆抗体

英文名称: Anti-ZFAND5 rabbit polyclonal antibody

别 名: ZA20D2; ZNF216; ZFAND5A

抗 原: ZFAND5

储 存: 冷冻(-20℃)

宿 主: Rabbit

相关类别: 一抗

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

Involved in protein degradation via the ubiquitin-prot easome system. May act by anchoring ubiquitinated proteins to the proteasome. Plays a role in ubiquitin-mediated protein degradation during muscle atrophy. Plays a role in the regulation of NF-kappa-B activation and apoptosis. Inhibits NF-kappa-B activation trigg ered by overexpression of RIPK1 and TRAF6 but not of RELA. Inhibits also tumor necrosis factor (TNF), IL-1 and TLR4-induced NF-kappa-B activation in a dose -dependent manner. Overexpression sensitizes cells to TNF-induced apoptosis. Is a potent inhibitory factor for osteoclast differentiation.



Applications:	ELISA, WB, IHC
Name of antibody:	ZFAND5
Immunogen:	Full length fusion protein
Full name:	zinc finger, AN1-type domain 5
Synonyms:	ZA20D2; ZNF216; ZFAND5A
SwissProt:	O76080
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human thyroid cancer and human lung cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	23 kDa
WB Positive control:	Mouse fat tissue lysate
WB Recommended dilution:	500-2000

