

兔抗 ACKR2 多克隆抗体

中文名称：兔抗 ACKR2 多克隆抗体

英文名称：Anti-ACKR2 rabbit polyclonal antibody

别名：atypical chemokine receptor 2; D6; hD6; CCR9; CCBP2; CCR10; CMKBR9

相关类别：一抗

抗原：ACKR2

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes a beta chemokine receptor, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptor-mediated signal transduction are critical for the recruitment of effector immune cells to the inflammation site. This gene is expressed in a range of tissues and hemopoietic cells. The expression of this receptor in lymphatic endothelial cells and overexpression in vascular tumors suggested its function in chemokine-driven recirculation of leukocytes and possible chemokine effects on the developme

| | |
|------------------------------------|--|
| | nt and growth of vascular tumors. This receptor appears to bind the majority of beta-chemokine family members; however, its specific function remains unknown. This gene is mapped to chromosome 3p21.3, a region that includes a cluster of chemokine receptor genes. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | ACKR2 |
| Immunogen: | Synthetic peptide of human ACKR2 |
| Full name: | atypical chemokine receptor 2 |
| Synonyms: | D6; hD6; CCR9; CCBP2; CCR10; CMKBR9 |
| SwissProt: | O00590 |
| ELISA Recommended dilution: | 5000-10000 |
| WB Predicted band size: | 43 kDa |
| WB Positive control: | Human placenta tissue lysate |
| WB Recommended dilution: | 200-1000 |



