

兔抗 BLNK 多克隆抗体

- 中文名称：兔抗 BLNK 多克隆抗体
- 英文名称：Anti-BLNK rabbit polyclonal antibody
- 别名：B-cell linker; bca; AGM4; BASH; LY57; SLP65; BLNK-S; SLP-65
- 抗原：BLNK
- 相关类别：一抗
- 储存：冷冻 (-20℃)
- 宿主：Rabbit
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphobla

	stic leukemia. Alternatively spliced transcript variants have been found for this gene.
Applications:	ELISA, IHC
Name of antibody:	BLNK
Immunogen:	Fusion protein of human BLNK
Full name:	B-cell linker
Synonyms:	bca; AGM4; BASH; LY57; SLP65; BLNK-S; SLP-65
SwissProt:	Q8WV28
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and human gastric cancer
IHC Recommend dilution:	50-300



